



Okeanos Explorer ROV Dive Summary

Dive Information	
Dive Map	
Site Name	Hadal Trough "Kinono"
Expedition Coordinator(s)	Brian RC Kennedy, Nick Pawlenko
ROV Lead(s)	Karl McLetchie
Science Team Lead(s)	Amanda Demopoulos and Steven Auscavitch
General Area Descriptor	Phoenix Islands Protected Area
ROV Dive Name	
Cruise	EX-17-03


	Asako Matsumoto	Chiba Institute of Technology (Chitech),	amatsu@gorgonian.jp
	Benjamin Frable	Scripps Institution of Oceanography	bfrable@ucsd.edu
	Bruce Mundy	NOAA NMFS Pacific Islands Fisheries Science Center	bruce.mundy@noaa.gov
	Christopher Kelley	University of Hawaii	ckelley@hawaii.edu
	Jaymes Awbrey	University of Louisiana, Lafayette	jawbrey@louisiana.edu
	Jill Bourque	US Geological Survey Wetland and Aquatic Research Center	jbourque@usgs.gov
	Les Watling	University of Hawaii at Manoa	watling@hawaii.edu
	Natalie Summers	University of Hawaii at Manoa	nsummers@hawaii.edu
	Nolan Barrett	FAU Harbor Branch Oceanographic Institute	barrettnh@g.cofc.edu
	Peter Auster	Mystic Aquarium & UConn	peter.auster@uconn.edu
	Randi Rotjan	Boston University	rrotjan@bu.edu
Santiago Herrera	Lehigh University	sherrera@alum.mit.edu	

	Scott France	University of Louisiana at Lafayette	france@louisiana.edu
	Steve Auscavitch	Temple University	steven.auscavitch@temple.edu
	Taylor Heyl	WHOI	theyl@whoi.edu
	Timothy Shank	Woods Hole Oceanographic Institution	tshank@whoi.edu
	Tina Molodtsova	P.P.Shirshov Institute of Oceanology RAS	tina.molodtsova@gmail.com
	Witting Jan	Sea Education Association	jwitting@sea.edu
	Brendan Roak	Texas A&M University	broark@geos.tamu.edu
Purpose of the Dive	<p>The general goal of this dive is to acquire baseline information on deep sea habitats, seafloor geology, and biological communities at hadal depths in the Phoenix Islands Protected Area (PIPA). We will be diving in one of the few non-trench associated hadal zones in the Pacific Ocean. Most hadal environments are virtually unexplored and there is little understanding of what resources might be protected at such depths. This dive will provide some perspective on biological resources (e.g. fishes, biogenic structures) as well as seafloor geology at these depths. The geological processes for formation of the Nova-Canton Trough could also be associated with the formation of this hadal trough in the Phoenix Islands.</p>		
Description of the Dive	<p>EX1703 dive # 16 was our deepest dive for the expedition, descending to 5862m in a deep trough feature, southwest of the Phoenix Islands. The seafloor was composed of scattered and emplaced rocks coated in manganese iron oxide crust, with brown sediment drape occurring in patches. The sediment appeared more organic rich and darker in color than on our previous dives in the region, speckled with calcareous foram tests. This dive should be nicknamed “white house, black market”. Throughout the dive, both on the steep slope and sedimented ledge at the end of the dive, we encountered taxa that lacked pigmentation, which contrasted with the dark color of the seafloor. We observed white anemones, holothurians, brisingids (<i>Freyastera</i>- some with swollen arms filled with gonads), cladorhizid sponges (2 species),</p>		


comatulid crinoids with eggs, solitary hydroid, stalked sponges with amphipod associates (Stegocephalidae), slit shell gastropod (Anatomidae: *Anatoma* sp.), sea pigs (unknown Elpidiidae), pentagonal ophiuroid, and a branching bryozoan. We also saw a large amphipod (e.g., Epimeriidae), red caridean shrimp, carnivorous tunicates, large pink mysid shrimps, fan-shaped xenophyophores, brown globe forams (cf. Komokioidea), long, mud tube with a white polychaete inside (cf. Onuphidae), and a cusk eel (Ophidiidae: *Alcockia rostrata* or *Bathyonus caudalis*).

We observed some unexpected fauna, including priapulids feeding on the sediment surface, large purple holothurians (*Psychropotes*), and a rare monoplacophoran mollusc. At the end of the dive, along a sedimented plateau composed of hard pavement, we saw a white seapen (cf. *Umbellula*) and collected the only black coral (cf. *Abyssopathes* sp.) observed on the dive, at ~ 5775 m.

Overall Map of the ROV Dive Area



Close-up Map of Main Dive Site



Representative Photos of the Dive



The seafloor was composed of scattered and emplaced rocks coated in manganese iron oxide crust, with brown sediment drape occurring in patches.

Possibly a monoplacophoran mollusc

Samples Collected

Sample

Sample ID	EX1703_20170323T214437_D2_DIVE16_SPEC01GEO	
Date (UTC)	20170323	
Time (UTC)	21:44:37	
Depth (m)	5859.54	
Temperature (°C)	1.37	
Field ID(s)	Mn-Coated Rock	
Comments		

Sample

Sample ID	EX1703_20170324T010307_D2_DIVE16_SPEC02BIO	
Date (UTC)	20170324	
Time (UTC)	01:03:07	
Depth (m)	5772.36	
Temperature (°C)	1.35	

Field ID(s)	Antipatharia	
Comments		

Please direct inquiries to:

NOAA Office of Ocean Exploration & Research
1315 East-West Highway (SSMC3 10th Floor)
Silver Spring, MD 20910
(301) 734-1014