

Okeanos Explorer ROV Dive Summary

Dive Information			
General Location	Ove 3: Beethoven Ridge Ove 19: Mendelsohn Seamount Dive 19: Mendelsohn Seamount Dive 2: Berich Ridge Ove 20: Middle bank		
General Area Descriptor	Just outside of PMNM, US EEZ Hawaii		
Site Name	Middle Bank		
Science Team Leads	John R. Smith/Meagan Putts		
Expedition Coordinator	Kasey Cantwell		
ROV Dive Supervisor	Karl McLetchie		
Mapping Lead	Mike White		
ROV Dive Name			
Cruise	EX1708		
Leg	-		
Dive Number	DIVE20		
Equipment Deploye	d		
ROV	Deep Discoverer		
Camera Platform	Seirios		
ROV Measurements	🖂 СТD	🔀 Depth	Altitude

	Scanning Sonal	r	🛛 USBL Posi	tion	Heading
	Pitch		🔀 Roll		HD Camera 1
	🛛 HD Camera 2		Low Res C	am 1	Low Res Cam 2
	🛛 Low Res Cam 3	}	Low Res C	Cam 4	Low Res Cam 5
Equipment Malfunctions			·		
	Dive Summary: EX1708_DIVE20				
ROV Dive Summary (from processed ROV data)	In Water: 2017-09-26T18:26 22°, 45.217' N ; 16		9-26T18:26:12	198000	·
	Out Water: 2017-09-27T02:23:51.628 22°, 45.330' N ; 160°, 55.8				
	Off Bottom: 2017-09-27T02:00:25.761000 22°, 45.281' N ; 160°, 55.957' W				
	On Bottom:	om: 2017-09-26T18:57:15.580000 22°, 45.249' N ; 160°, 55.692' W			
	Dive duration: 7:57:39				
	Bottom Time: 7:3:10				
	Max. depth: 477.7 m		m		
Special Notes					
	Name	Email		Affiliation	
Scientists Involved (please provide name, location, affiliation, email)	Asako Matsumoto	amatsu@gorg	onian.jp	· · · ·	ploration Research a Institute of Technology
	Bruce Mundy	bruce.mundy@noaa.gov		NOAA NMFS Pacific Islands Fisheries Science Center	
	Dhugal Lindsay	dhugal@jamstec.go.jp JAMSTEC			
	Frank Parrish	Frank.Parrish@	pnoaa.gov	PIFSC	
	John Smith	jrsmith@hawa	iii.edu	University of Hawaii	
	Katie Wagner	katie.wagner@	onoaa.gov	NOAA OER	
	Kevin Kocot	kmkocot@ua.	edu	The Universi	ty of Alabama
	Les Watling	watling@hawa	aii.edu	University of	⁻ Hawaii at Manoa
	Meagan Putts	Meagan.putts	@noaa.gov	University of	
	Megan McCuller	mccullermi@g	mail.com	Williams-My Program	stic Maritime Studies



	Michael Parke	michael.parke@noaa.gov	NOAA PIFSC
			FAU Harbor Branch Oceanographic
	Nolan Barrett	barrettnh@g.cofc.edu	Institute
	Scott France	france@louisiana.edu	University of Louisiana at Lafayette
	Tina Molodtsova	tina@ocean.ru; tina.molodtsova@gmail.com	P.P.Shirshov Institute of Oceanology RAS
Purpose of the Dive	This dive, located just outside the boundary of PMNM, was carried out at the request of the Monument staff and NOAA NOS NCCOS. The objectives were 1) to explore fish and invertebrate boundary communities and 2) ground truth coral habitat suitability and taxonomic richness models based on multibeam sonar bathymetry and slope data. Management unit species within the precious coral and bottomfish fisheries were of particular interest. The selection of this dive site on Middle Bank was based on high-resolution multibeam bathymetry, slope, and backscatter data revealing the presence of a pinnacle feature that likely had populations of bottomfish and precious corals. In addition, regional coral habitat suitability models indicated it was close to an area that had high potential habitat suitability for the occurrence of seven or more genera of coral species. Draft models were produced as a collaborative effort by NOAA NOS NCCOS as part of a BOEM-funded Biogeographic Assessment of the Main Hawaiian Islands. More information about the project can be found at:		
Description of the Dive	massive amounts of dead deep-sea coral frees collecting at the base of the cone and		
	depth near the bas surface, only intern <i>mitsukurii</i>) and cor close proximity. A at 476 m. D2 bega composed of carbo <i>Hemicorallium</i> sp. <i>Acanthogorgia</i> sp. an long dead gold	se of the cone. The seafloor we rupted by thin carbonate-like I rals (Primnoidae, <i>Ennalopsamr</i> gentle slope collecting a large in imaging "dissolution holes" onate. The first documented p pink coral, was observed at tir coral, followed by an Asteropol coral branch at time 19:18 (47	ottom at time 16:57 at 475 m water as a flat and featureless brown hard ayers. Two Dogfish sharks (<i>Squalus</i> <i>nia</i> sp. "purple') were observed in amount of dead coral trees was seen in the pavement surface suggesting it is precious coral on Middle Bank, a me 19:14 (475 m) underneath a large hiura sp. brittle star perched on top of 5 m). Darker outcrops and ledges came n at 471 m. The dark matter also




surrounded (flowed around?) some rugose and lighter colored (tan) outcrops likely to be fossil coral reef. The slope increased from flat to 20-30°, which remained generally consistent for most of the dive until D2 rose closer to the summit. More inclined outcropping slabs with dissolution beneath came into view at 470 m. The depth of 470-471 meters may mark the transition between all carbonate and mixed mode substrate, at least on this one dive track. A suspended ledge "tongue" with many corals on the tip was observed at time 19:47 and Codling, Laemonema rhodochir at time 19:54. Closeups of the dark material forming a ledge showed a tan colored matrix speckled with shiny black bits -all highly confusing. To add to that, a thin vein of tan material persisting between layers of dark material on an incline were observed at time 20:12. To help alleviate the confusion, a wide view of a carbonate tongue (tan) surrounded by tuff (volcaniclastic, dark) was captured at time 20:20 (455 m). A close-up a monkfish, Lophiodes micanthus, was obtained at time 20:20 (455 m). More contacts with tongues of dark material flowing through and around tan substrate, with numerous gold, Kulamanamana haumeaae, and bamboo corals, Acanella sp. and Orstomisis sp., at 449 m. A Plinthaster sp. cookie star was imaged at time 20:31 (448 m). No available rocks had been seen until a collection of a lone angular talus piece from the open and featureless slope was collected at time 20:44 (447 m). Later examination aboard confirmed it was of volcanic origin, likely basalt, although highly altered and brown in color outside. It was split open along an existing crack and the matrix and numerous clasts were found to be reddish-orange with no vesicles. Gold corals abound in this location with the present current flow from the north at ~0.5 knot. An example of the "Midas" growth stage of a bamboo coral being recolonized by a gold coral was imaged at time 20:52 (443 m). An orange-red Leiopathes sp. coral with two nearby squat lobsters were seen at time 20:57 (439 m) and two specimens of black coral were collected at 405 m. A large school of Amberjacks, Seriola durmerili, were first seen at 397 m and likely the same school persisted throughout the remainder of the dive. The slope increased to ~45° at 378 m with the same alternating tan and dark substrate. A pencil urchin, Stereocidaris hawaiiensis; scleractinian coral, Equchipsammia sp.; and a colorful squat lobster, Eummunida sp., were observed between times 23:18 to 23:39 and a wiry white plexaurid coral was collected at 23:33 (374 m). A close-up of a gold coral base showed on-lapping of the dark (volcaniclastic?) material, indicating that the eruption events were coeval with at least the older corals. This may explain the large number of dead gold coral bases found on the slope and at the base of this cone. A Chaunax umbrinus angler fish with bright orange and yellow camouflage pattern along with a number of small sponges were imaged at time 00:00 (359 m). A Bandfish, Owstonia hawaiiensis, was imaged at time 00:06 (356 m), apparently somewhat unusual in these environs. A nice wide view of the tan material over dark was had at time 00:07 (352 m). Some shimmering water had been observed and the temperature varied by ~1° during these episodes, this time at 00:18 (332 m). It may be explained by water mass mixing at these relatively shallow depths as no geologic, hydrologic, or hydrothermal activity is thought to presently be active on Middle Bank. Interestingly, the pilots reported that the ambient temperature dropped when the shimmering was observed. Possible cooling fractures in the volcanic (dark) material were observed running up and downslope at time 00:25 (335 m). Possibly a new fish, orange with big black eyes, that stumped the experts was imaged at time 00:30 (330 m). The assemblage of black coral trees was noted to be getting denser and larger as the summit was approached. At time 00:45, D2 came upon vertical walls of massive outcrop with densely packed black corals lining all edges of the rocks composed of the tan material. Another raised vertical ledge of tan material covered with many black corals was traversed, along with a derelict piece of fishing line, at time 00:52 (312 m). Another local high, a false summit ridge, was ascended at time 01:04 (307 m). The substrate



soon after became confusing again, looking like volcanic tuff but not as dark as before, now in between the dark and tan material – perhaps brown. An Antipathella sp.? coral was collected at time 01:24 (304 m) since it is by far the dominate species on the upper portion of this cone feature. A "conger-gation" of nine large eels, Conger oligoporus, peering out of a gap in a large outcrop was seen at time 01:33 (301 m). D2 descended into a gully between massive outcroppings of brown material at time 01:42, and after rising to the top a "living rock" of anemones, demosponges, corallimorpharians, etc. covering its entire observable surface was documented at time 01:47 (295 m). The anemones were also seen to carpet the tops of nearby outcrops into the distance carpeting the rock face, mimicking the look of Mn-crust, even resembling basalt with vesicles at a distance! Undulating surfaces of massive outcrops with fractures, gulleys, and isolated blocks were observed on the last run along the summit at time 01:53. D2 left bottom at time 02:00 from a water depth of 296 m as a large school of small fish quickly swam underneath the ROV.

Overall Map of the ROV Dive Area



Representative Photos of the Dive





Curious green-eyed dogfish, <i>Squalus mitsukurii</i> , swiftly swimming by D2		A before and after shot of colonization by gold coral (left), Kulamanamana haumeaae, and likely host colony species, bamboo coral (right), <i>Acanella</i> sp.
Large school of A boldly circling D2	mberjacks, Seriola durmerili,	Large boulder 100% covered with demosponges, anemones, corallimorpharians and more!
Samples Collecte	d	
Sample		
Sample ID	EX1708_D2_DIVE20_SPEC01GE O	
Date (UTC)	9/26/2017	
Time (UTC)	20:45	
Depth (m)	447.7	
Temperature (°C)	8.8	
Field ID(s)	Angular basalt talus from open slope. Brown in color. Split open along existing crack to reveal that the matrix and numerous clasts are reddish- orange in color with no vesicles.	
Commensal ID and Field Identification		 A02 Poriferan A03 <i>Hemicorallium</i> sp. A04 Scleractinia Single Polyp 1 A05 Scleractinia Colonial Polyps 2 A06 Scleractinia Colonial Polyps 3 Dead A07 Hydrozoan
Comments		





Sample		
Sample ID	EX1708_D2_DIVE20_SPEC02BIO	SANNE GY SPACE
Date (UTC)	9/26/2017	
Time (UTC)	22:27	
Depth (m)	405.1	
Temperature (°C)	9.9	
Field ID(s)	Myriopathes sp.	
Commensal ID and Field Identification	EX1708_D2_DIVE20_SPEC02BIO_4	A01 Scalpellidae
Comments		
Sample		
Sample ID	EX1708_D2_DIVE20_SPEC03BIO	
Date (UTC)	9/26/2017	
Time (UTC)	22:33	
Depth (m)	404.2	and a series
Temperature (°C)	10.1	
Field ID(s)	Myriopathes ulex?	
Commensal ID and Field Identification		
Comments		
Sample		
Sample ID	EX1708_D2_DIVE20_SPEC04BIO	
Date (UTC)	9/26/2017	
Time (UTC)	23:33	
Depth (m)	374.4	
Temperature (°C)	10.5	
Field ID(s)	Plexauridae	
Commensal ID and Field Identification	EX1708_D2_DIVE20_SPEC04BIO_4	A01 Actinarian
Comments		



Sample		
Sample ID	EX1708_D2_DIVE20_SPEC05BIO	
Date (UTC)	9/27/2017	
Time (UTC)	01:24	
Depth (m)	305.0	
Temperature (°C)	12.3	
Field ID(s)	Antipathella sp.?	
Commensal ID and Field Identification	EX1708_D2_DIVE20_SPEC05BIO_A	01 Actinarian
Comments		

Please direct inquiries to:

NOAA Office of Ocean Exploration & Research 1315 East-West Highway (SSMC3 10th Floor) Silver Spring, MD 20910 (301) 734-1014

