




Okeanos Explorer ROV Dive Summary

Dive Information	
General Location	
General Area Descriptor	Blake Plateau
Site Name	Richardson Ridge
Science Team Leads	Leslie Sautter / Cheryl Morrison
Expedition Coordinator	Kasey Cantwell
ROV Dive Supervisor	Bobby Mohr
Mapping Lead	Derek Sowers
ROV Dive Name	
Cruise	EX1806
Leg	-
Dive Number	DIVE07
Equipment Deployed	
ROV	Deep Discoverer

Christian Jones	NMFS	christian.jones@noaa.gov
	Dept of Invertebrate Zoology, NMNH	
Christopher Mah	Smithsonian	brisinga@gmail.com
Derek Sowers	OER	derek.sowers@noaa.gov
Enrique Salgado	NCCOS	enrique.salgado@noaa.gov
Erik Cordes	Temple University	ecordes@temple.edu
Ervan Garrison	University of Georgia	egarriso@uga.edu
	University of South Florida St. Petersburg	
Heather Judkins		Judkins@mail.usf.edu
	University of the Azores and Senckenberg am Meer, Germany	
Íris Sampaio		irisfs@gmail.com
		jlunden@temple.edu; jlunden@haverford.edu
James Lunden	Temple University	
	NOAA OER - Hawaii Sea Grant Knauss Fellow	
James Murphy		james.murphy@noaa.gov
Jill Bourque	US Geological Survey	jbourque@usgs.gov
	University of Bergen (Norway)	
Joana Xavier		joanarxavier@gmail.com
	Harbor Branch Oceanographic Institute	
John Reed		jreed12@fau.edu
Kate Rose	NOAA NCEI	kate.rose@noaa.gov
Kenneth Sulak	USGS (ret.)	ksulak@usgs.gov
Kevin Jerram	UNH	kjerram@ccom.unh.edu
	Univeristy of Louisiana, Lafayette	
Lauren Walling		c00305146@louisiana.edu
	University of Hawaii at Manoa	
Les Watling		watling@hawaii.edu
Leslie Sautter	College of Charleston	Sautterl@cofc.edu
	North Carolina Museum of Natural Sciences	
Megan McCuller		mccullermi@gmail.com
Michael Vecchione	NOAA/NMFS National Systematics Lab	vecchiom@si.edu
Mike Ford	NOAA Fisheries	michael.ford@noaa.gov
Nolan Barrett	South Carolina University	barrettnh@g.cofc.edu
Rachel Bassett	NOAA NCCOS DCEL	rachel.bassett@noaa.gov





	Robert Carney	Louisiana State Univ	rcarne1@lsu.edu
	Sandra Brooke	Florida State University	sbrooke@fsu.edu
	Santiago Herrera	Lehigh University	sherrera@alum.mit.edu, sah516@lehigh.edu
	Scott Allen	NOAA Ship Okeanos Explorer	
	Scott France	University of Louisiana at Lafayette	france@louisiana.edu
	Scott Harris	College of Charleston	harriss@cofc.edu
	Sophie Alpert	College of Charleston	alpertsl@g.cofc.edu
	Stephanie Bush	Smithsonian	stephalopod@gmail.com
	Tara Harmer		
	Luke	Stockton University	luket@stockton.edu
	Tina Molodtsova	Shirshov Institute of Oceanology RAS	tina@ocean.ru; tina.molodtsova@gmail.com
	Treyson Gillespie	College of Charleston	gillespieta@g.cofc.edu
	Victoria Gitto	College of Charleston	geogittotm@gmail.com
	Zach Proux	College of Charleston	prouxzs@g.cofc.edu
Purpose of the Dive	<p>This dive site was selected for its knife-like ridge at the top of a steep 100 m relief slope that had gradients exceeding 30°. Its position on the outer edge of the Gulf Stream's flow raised the question if it was of biogenic (i.e., coral rubble mound) or geologic (erosional scarp of rock layers) origin. The previous 2 dives explored mound features, revealing origin from massive accumulation of stoney coral skeletons. All 4 mounds explored hosted dense living coral and sponge communities. The Richardson Ridge was explored to compare biological communities on the ridge's slope (in the lee of the Gulf Stream) to those on the ridge crest, and to compare these areas with those of the previous 2 dives. This dive was part of a series that investigates the similarities and differences in community composition between deepwater habitats of the SE US continental margin. This site is within a large under-explored area of the Miami Stetson CHAPC and is a high priority region for the Deep Search project, with the primary objective to identify presence / absence of deepwater corals and benthic communities in areas with the potential for offshore development.</p> <p>This area was first mapped during EX-18-05. New information will inform biogeographic patterns in the region, critical for refining habitat prediction models. Diving in the area will provide important information to groundtruth these models.</p>		




<p>Description of the Dive</p>	<p>The dive ascended the steep slope (>30o at times) of what turned out to be an eroded coral mound. The steep slopes were covered with coral rubble, and as the ROV ascended, more coral communities were observed. The crest of the ridge was covered with 1 to 1.5 m high mounds of <i>Lophelia pertusa</i> and <i>Madrapora oculata</i>. Eunicid polychaetes were observed in the coral matrix of both framework-building corals. The octocoral <i>Plumarella</i> was common, and several colonies of the bamboo coral, <i>Keratoisis</i>, and Anthothelidae were observed. Soft corals (Alcyonacea: <i>Anthomastus</i> and another Nephtheidae) were also common. Several individuals of a hormathiidae anemone (Actiniaria) were observed. Squat lobsters from the Galattheoidea and Chirostyloidea were observed, along with hermit crabs and the golden crab <i>Chaceon fenneri</i>. The large red shrimp <i>Pleisopenaeus armatus</i> was observed on coral rubble. Echinoderms included: the Echinothuroid <i>Aerosoma</i> (pancake urchin), brittle stars (possibly <i>Ophicantha bidentata</i>) in coral rubble, sea stars (Asteroidea) including <i>Gilbertaster caribbaea</i> observed feeding on the octocoral <i>Plumarella</i> sp. and <i>Chondraster grandis</i> (Poraniidae), Echinus and Cidaroid Pencil urchins. The hexactinellid sponges <i>Aphrocallistes beatrix</i> (with yellow zoanthids) and <i>Vazella pourtalesi</i> were seen, along with <i>Geodia</i> sp. (demosponge). The most common fish was <i>Synaphobranchus kaupii</i>. Other fishes observed included a skate (<i>Fenestraja plutonia</i>), an anglerfish (<i>Lophioides beroe</i>), the coral hake <i>Laemonema melanurum</i> and a goosefish, <i>Lophiodes beroe</i>.</p> <p>The current-facing slope was more densely populated with live <i>Lophelia</i> corals as compared to the lee-side slope. The narrowness of the ridge crest appears to be in part the result of erosional slumping on the lee-side. Such slumping over the life history of this mound would result in steep slopes and a narrow ridge crest, rather than the rounded mound morphology seen on earlier dives.</p>	
<p>Notable Observations</p>	<p>Learning that these elongate, steep ridge areas are actually coral mounds is a significant finding, as these features occur in many areas of the region in addition to the rounded mounds seen throughout Stetson Mesa. The amount of live coral was quite high compared to other coral mounds visited.</p> <p>Observation of the seastar <i>Gilbertaster caribbaea</i> feeding was a first! <i>Plumarella</i> appeared to be this species' food of choice, as we encountered three dining individuals.</p>	
<p>Community Presence/ Absence (<i>community is defined as more than two species</i>)</p>	<p><input checked="" type="checkbox"/> Corals and Sponges Present</p> <hr/> <p><input type="checkbox"/> Chemosynthetic Community Present</p> <hr/> <p><input type="checkbox"/> High biodiversity Community Present</p>	<p><input type="checkbox"/> Active Seep or Vent</p> <hr/> <p><input type="checkbox"/> Extinct Seep or Vent</p> <hr/> <p><input type="checkbox"/> Hydrates Present</p>
<p>Overall Map of the ROV Dive Area</p>		<p>Close-up Map of Main Dive Site</p>





Representative Photos of the Dive




At the beginning of the dive, the substrate was gently sloping and matted with dead *Lophelia* skeletal matter. A *Plesiopenaeus* sp. shrimp was observed here.

On the steeply sloped ascent we encountered larger mounds of skeletal framework, with living *Lophelia pertusa* and the octocoral *Plumarella* sp. A goosefish is also shown in the image.




The southwestward-facing slope (into the current) near the ridge crest was covered with large 1 m mounds of *Lophelia*.

At the ridge crest, large mounds of *Lophelia* and *Madrapora* were found.



The knife-like ridge is likely the result of erosional slumping, shown at right. Debris piles at the base and becomes habitat (left).



A view of both sides of the ridge show the increased abundance of live *Lophelia* on the current-facing side, at left.



Three different *Gilbertaster caribbaea* sea stars were each observed feeding on the octocoral *Plumarella* sp.



The seastar *Chondraster grandis* (Poraniidae) seen on the coral rubble.



Several golden crabs (*Chaceon fenneri*) were seen living within the coral mounds.



An oreo (*Neocyttus* sp) was observed over live *Lophelia*.



This type of crinoid, *Zenometra columnaris*, was observed in abundance, living on the coral skeletal framework.

Samples Collected

Sample

Sample ID	D2_DIVE07_SPEC01BIO	
Date (UTC)	20180621	
Time (UTC)	212230	
Depth (m)	787.54	
Temperature (°C)	8.31	
Field ID(s)	<i>Madrepora oculata</i>	

Reason for Collection	<i>Aspire target species</i>
-----------------------	------------------------------

Notes	
-------	--

Associates	Associate ID	Field Identification	Notes
	A01	Ophiuroidea	
	A02	Polychaeta	
	A03	Eunicidae	Include in ASPIRE

Sample

Sample ID	D2_DIVE07_SPEC02BIO	
Date (UTC)	20180621	
Time (UTC)	212634	
Depth (m)	788.25	
Temperature (°C)	8.29	
Field ID(s)	<i>Lophelia pertusa</i>	



Reason for Collection	<i>Aspire target species</i>		
Notes			
Associates	Associate ID	Field Identification	Notes
	A01	Decapoda	gravid female shrimp
	A02	Ophiuroidea	
	A03	Hydrozoa	hydroid
	A04	Hydrozoa	hydroid
	A05	Demospongiae	
	A06	Polychaeta	
	A07	Eunicidae	Include in ASPIRE
Sample			
Sample ID	D2_DIVE07_SPEC03GEO		
Date (UTC)	20180621		
Time (UTC)	220647		
Depth (m)	763.74		
Temperature (°C)	10.45		
Field ID(s)	Coral rubble		
Reason for Collection	<i>Site characterization</i>		
Notes			
Associates	Associate ID	Field Identification	Notes
	A01	Foraminifera	Agglutinated
	A02	Ophiuroidea	
	A03	Gastropoda	



Please direct inquiries to:

NOAA Office of Ocean Exploration & Research
 1315 East-West Highway (SSMC3 10th Floor)
 Silver Spring, MD 20910
 (301) 734-1014

