


Okeanos Explorer ROV Dive Summary

Dive Information



ROV Dive Name

Cruise	EX1903L2
Dive Number	Dive02

Equipment Deployed

Seirios ✓ CTD ✓ Scanning Sonar ✓ Pitch ✓ HD Camera 2 ✓ Low Res Cam 3 Dive Summary: EX1903	✓ Depth ✓ USBL Position ✓ Roll ✓ Low Res Cam 1 ✓ Low Res Cam 4	✓ Altitude ✓ Heading ✓ HD Camera 1 ✓ Low Res Cam 2 ✓ Low Res Cam 5	
✓ Scanning Sonar ✓ Pitch ✓ HD Camera 2 ✓ Low Res Cam 3 Dive Summary: EX1903	✓ USBL Position ✓ Roll ✓ Low Res Cam 1 ✓ Low Res Cam 4	✓ Heading✓ HD Camera 1✓ Low Res Cam 2	
✓ Pitch ✓ HD Camera 2 ✓ Low Res Cam 3 Dive Summary: EX1903	✓ Roll ✓ Low Res Cam 1 ✓ Low Res Cam 4	✓ HD Camera 1 ✓ Low Res Cam 2	
✓ HD Camera 2 ✓ Low Res Cam 3 Dive Summary: EX1903	✓ Low Res Cam 1 ✓ Low Res Cam 4	✓ Low Res Cam 2	
✓ Low Res Cam 3 Dive Summary: EX1903	✓ Low Res Cam 4		
Dive Summary: EX1903		✓ Low Res Cam 5	
•	L2_DIVE02		
•	L2_DIVE02		
^^^^^			
		^^^^	
In Water:	2019-06-22T12:48:41.424703		
29°, 6.6	541' N ; 79°, 26.701' W		
On Bottom:	2019-06-22T13:45:28.171631		
29°, 6.849' N ; 79°, 26.867' W			
Off Bottom:	2019-06-22T19:29:58.780109		
29°, 6.5	539' N ; 79°, 26.855' W		
Out Water:	2019-06-22T22:40:16.027195		
29°, 7.7	785' N ; 79°, 26.9' W		
Dive duration:	9:51:34		
Bottom Time:	5:44:30		
Max. depth:	784.0 m		
igh currents at the surfa t 12 kts from the west.	ice (~1.5 kts) slowed the descent	and acquiring the bottom. Wind were	
בו בו	29°, 6.6 On Bottom: 29°, 6.8 Off Bottom: 29°, 6.9 Out Water: 29°, 7.5 Oive duration: Bottom Time: Max. depth:	29°, 6.641' N; 79°, 26.701' W 2019-06-22T13:45:28.171631 29°, 6.849' N; 79°, 26.867' W 2019-06-22T19:29:58.780109 29°, 6.539' N; 79°, 26.855' W 2019-06-22T22:40:16.027195 29°, 7.785' N; 79°, 26.9' W 20ive duration: 9:51:34 3ottom Time: 5:44:30 Max. depth: 784.0 m	



Scientists Involved (provide name, affiliation, email)

Name	Affiliation	Email
Alexis Weinnig	Temple University	aweinnig@temple.edu
Allen Collins	NOAA's National Systematics Lab	Collinsa@si.edu
Amy Wagner	California State University, Sacramento	amy.wagner@csus.edu
Asako Matsumoto	Chiba Institute of Technology	amatsu@gorgonian.jp
Charles Messing	Nova Southeastern University	messingc@nova.edu
Christopher Mah	Dept. of Invertebrate Zoology, NMNH Smithsonian	brisinga@gmail.com
Danielle Power	NOAA Ship Okeanos Explorer	danielle.l.power@noaa.gov
Elizabeth Fraser	NOAA NCCOS	gugliottief@gmail.com
Elizabeth Gugliotti	NOAA Deep Sea coral ecology lab	elizabeth.gugliotti@noaa.gov
Enrique (Ren) Salgado	NOAA CSS	enrique.salgado@noaa.gov
Erik Cordes	Temple University	ecordes@temple.edu
Georgios Kazanidis	University of Edinburgh	georgios.kazanidis@ed.ac.uk
Herbert Leavitt	NOAA Office of Ocean Exploration and Research (OER Hollings Scholar)	herbert.leavitt@noaa.gov
J Dunn	NOAA OER	christopher.dunn@noaa.gov
John Reed	Harbor Branch Oceanographic Institute	jreed12@fau.edu
Kenneth Sulak	U.S. Geological Survey (Emeritus)	jumpingsturgeon@yahoo.com
Kevin Jerram	UNH	kjerram@ccom.unh.edu
Kevin Kocot	University of Alabama	kmkocot@ua.edu
Megan McCuller	North Carolina Museum of Natural Sciences	megan.mcculler@naturalsciences.or
Michael Vecchione	NOAA National Systematics Lab	vecchiom@si.edu
Mike Ford	NOAA	michael.ford@noaa.gov
Morgan Will	DCEL	morgan.will@noaa.gov
Nolan Barrett	Georgia Institute of Technology	barrettnh@g.cofc.edu
Shannon Hoy	NOAA OER	shannon.hoy@noaa.gov
Stacey Williams	Institute for Socio-Ecological Research	stcmwilliams@gmail.com




Stephanie Bush	Smithsonian Institution NMNH	stephalopod@gmail.com, bushsl@si.edu
Tara Harmer Luke	Stockton University	luket@stockton.edu, tara.luke@stockton.edu
Tina Molodtsova	P.P.Shirshov Institute of Oceanology RAS	tina@ocean.ru
Wilford Gardner	Texas A&M University	wgardner@ocean.tamu.edu

Dive Purpose	The three primary purposes of this dive were to: 1) explore underwater mounds in Million Mounds region that has high potential to be suitable habitat for deep sea coral and sponge communities, 2) investigate area with unusual backscatter data for ground-truthing, and 3) perform mid-water transects on the ascent to characterize and sample water column biota.
Dive Description	We started our dive a few hundred meters from the base of a cold-water coral mound at a depth of 770 m. Upon reaching the bottom, the ROV began transiting toward the Stetson Mesa South coral mound. We landed in an area of relatively soft sediment, however, a higher percentage of coral rubble vs. sand was observed. The coral rubble mixed with coarse-fine sands persisted throughout the majority of the dive and increased in abundance as we traversed up to the top of the mound and across the adjacent mounds. The general trend observed throughout the dive was a large abundance of coral rubble at the bottom of each mound with increasing abundance of live scleractinian coral coverage on the east to southeastern side of each mound. As we traversed along the base of the first mound, there was a substantial diversity of biota, including brittle stars, skates, sponges, pancake urchins, black coral and octocoral. There was a particularly high abundance of crinoids, particularly <i>Pentametrocrinus atlanticus</i> , the only feather star in the western Atlantic with five undivided arms. One individual <i>Pentametrocrinus</i> was observed swimming and it was noted that this was the first time this species was observed
	swimming through the water column. There was also a high abundance of cutthroat eels (Synaphobranchus) in the area in addition to three to four individual gulper sharks observed.
	As we proceeded up the first mound, the abundance of scleractinian coral rubble greatly increased and more live scleractinian (i.e. <i>Madrepora</i> , <i>Lophelia</i>) and octocorals were observed. Along the second mound that we explored, the occurrence and diversity of black coral species (i.e. <i>Leipoathes</i> , <i>Bathypathes</i> , <i>Stichopathes</i> , <i>Chrysopathes</i>) increased.
	After the benthic portion of the dive we conducted our first midwater exploration during this cruise. We completed transects at 700, 530, 500, and 300 meters and were able to complete three suction samples of pelagic organisms. The Deep Scattering Layer (DSL) was centered on 530 meters, this was added to the standard transect depths of 300, 500, and 700 for this dive. Both physonect and calycophoran siphonophores were observed during these transects, a typical occurrence for our midwater dives. We were able to collect a species of medusa in the genus <i>Haliscera</i> as our first suction sample for midwater exploration. Our observation of <i>Aegina citrea</i> will add to the records of geographic ranges and environmental conditions for this species.
Notable Observations	Extremely large abundance of crinoids at this site, dominated by <i>Pentametrocrinus atlanticus</i> . Video footage of swimming <i>Pentametrocrinus</i> - only second record of this genus swimming. Three to four possible gulper sharks swimming around the ROV.




Community	X Corals and Sponges
Presence/	✓ Chemosynthetic Community
Absence	X High biodiversity Community
(community is defined as more	✓ Active Seep or Vent
than two	✓ Extinct Seep or Vent
species)	✓ Hydrates
	,
Feature Type	Deepwater/coldwater stony coral reef (cold-water coral mounds)
SeaTube	https://data.oceannetworks.ca/SeaTubeV2?resourceTypeId=1000&resourceId=
(annotations	23621&diveId=2410
program) link	

Overall Map of the ROV Dive Area




EX1903L2 DIVE 02 - TRANSECT PLAN

(DRAFT)






Close-up Map of Main Dive Site



Benthic Portion of the dive



Midwater transects shown in 3D.



Representative Photos of the Dive




High coverage of dead coral rubble on and around mound feature.






Five arm crinoid (*Pentametrocrinus atlanticus*) on coral rubble. These were abundant throughout the beginning of the dive and one specimen was collected.



Slope of a cold-water coral mound with dead coral rubble and live white Plexaurid octocorals.





The jellyfish, Aegina citrea, in the water column observed during midwater exploration.

Samples Collected



Sample ID EX1903L2_D02_01B



Date (UTC)	20190622	
Time (UTC)	161741	
Depth (m)	749.0	
Temp. (°C)	8.267	
Field ID(s)	Crinoid	
Associates		
	Associates Sample ID	Field Identification
	No Associates	
Comments		



Sample ID	EX1903L2_D02_02B
Date (UTC)	20190622
Time (UTC)	182609
Depth (m)	740.2
Temp. (°C)	8.286
Field ID(s)	Plexaurid



Associates		
	Associates Sample ID	Field Identification
	EX1903L2_D02_02B_A01	Ophiuroidea
	EX1903L2_D02_02B_A02	Cirripedia
	EX1903L2_D02_02B_A03	Lophelia pertusa skeleton
	EX1903L2_D02_02B_A04	Madrepora oculata skeleton
Comments		



Sample ID	EX1903L2_D02_03B	
Date (UTC)	20190622	
Time (UTC)	195649	
Depth (m)	699.2	
Temp. (°C)	9.944	
Field ID(s)	Haliscera sp.	
Associates		
	Associates Sample ID	Field Identification
	No Associates	
Comments		





Sample ID	EX1903L2_D02_04B	
Date (UTC)	20190622	
Time (UTC)	204943	
Depth (m)	500.7	
Temp. (°C)	14.554	
Field ID(s)	Arctapodema sp	
Associates		
	Associates Sample ID	Field Identification
	No Associates	
Comments		





Sample ID	EX1903L2_D02_05B			
Date (UTC)	20190622	20190622		
Time (UTC)	205509			
Depth (m)	503.0			
Temp. (°C)	14.725			
Field ID(s)	Bathocyroe			
Associates				
	Associates Sample ID	Field Identification		
	No Associates			
Comments				

Please direct inquiries to:

NOAA Office of Ocean Exploration & Research 1315 East-West Highway (SSMC3 10th Floor) Silver Spring, MD 20910 (301) 734-1014

